

J003 REGISTRATION

24 JAN 2002

FILING BY "EXPRESS MAIL" UNDER 37 CFR 1.10

EK902078784US

"Express Mail" label mailing number

1/24/02

Date of Deposit

Form PTO-1590
(REV 10-95)

U. S. Department of Commerce Patent and Trademark Office

ATTORNEY'S DOCKET NUMBER

HF/5-22053/A/PCT

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)

10/048045

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

INTERNATIONAL APPLICATION NO.
PCT/EP 00/06934INTERNATIONAL FILING DATE
July 20, 2000PRIORITY DATE CLAIMED
July 28, 1999

TITLE OF INVENTION

Water-soluble granules of salen-type manganese complexes

APPLICANT(S) FOR DO/EO/US


Menno Hazenkamp, Frank Bachmann, Cornelia Makowka, Petr Kvita, Rolf Kuratli and Anita Schmidlin

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39 (1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau. (See attached Form PCT/IB/308)
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English 35 U.S.C. 371(c)(2).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☒ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☒ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included.

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ **A FIRST preliminary amendment.**
 - ☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: (See attached Form PCT/ISA/210)

U.S. APPLICATION NO. (if known, see 37 CFR 1.5) 10/048045		INTERNATIONAL APPLICATION NO. PCT/EP 00/06934		ATTORNEY'S DOCKET NUMBER HF/5-22053/A/PCT	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a) (1)-(5)):				CALCULATIONS PTO USE ONLY	
Search Report has been prepared by the EPO or JPO \$890.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) \$710.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)). \$740.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO. \$1040.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4). \$100.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$890.00	
Surcharge of \$130.00 for furnishing the oath of declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)). \$					
CLAIMS		NUMBER FILED	NUMBER EXTRA	RATE	
Total claims		15 - 20 =	0	X \$18.00 \$	
Independent claims		1 - 3 =	0	X \$84.00 \$	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)				+ \$280.00 \$	
TOTAL OF ABOVE CALCULATIONS =				\$890.00	
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).				\$	
SUBTOTAL =				\$890.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)). \$					
TOTAL NATIONAL FEE =				\$890.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$	
TOTAL FEES ENCLOSED =				\$	
				Amount to be: refunded	\$
				charged	\$890.00
a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. 03-1935 in the amount of \$890.00 to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 03-1935. A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
PLEASE ASSOCIATE THE ATTACHED APPLICATION WITH CUSTOMER NUMBER 000324 AND SEND ALL CORRESPONDENCE TO: JoAnn Villamizar, Ciba Specialty Chemicals Corporation Patent Department 540 White Plains Road P.O. Box 2005 Tarrytown, NY 10591-9005 DATE: JAN 24 2002					
				 SIGNATURE	
				Kevin T. Mansfield NAME Reg. No. 31,635	

CASE HF/5-22053/A/PCT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN RE PCT NATIONAL STAGE APPLICATION OF
MENNO HAZENKAMP ET AL
INTERNATIONAL APPLICATION NO. PCT/EP 00/06934
FILED: JULY 20, 2000
FOR: WATER-SOLUBLE GRANULES OF SALEN-
TYPE MANGANESE COMPLEXES
U.S. APPLICATION NO: UNASSIGNED
35 USC 371 DATE:

Group Art Unit:

Examiner:

Assistant Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Sir:

Kindly amend this application as follows prior to calculation of the filing fee and consideration on the merits.

IN THE CLAIMS



Please cancel claims 1-15.

Please add claims 16-30.


--16. **(new)** Water-soluble granules of salen-type manganese complexes, comprising

- a) from 1 to 89 % by weight of a water-soluble salen-type manganese complex,
- b) from 10 to 95 % by weight of a dissolution restrainer,
- c) from 0 to 20 % by weight of a further additive and
- d) from 1 to 15 % by weight of water, based on the total weight of the granules.

17. **(new)** Granules according to claim 16 that comprise as manganese complex a compound of formula



or



wherein

A is an anion;

m, n and p are each independently of the others 0, 1, 2 or 3,


R₄ is hydrogen or linear or branched C₁-C₄alkyl,

Y is a linear or branched alkylene radical of formula $-\text{C}(\text{R}_4)_2\text{-}$, wherein r is an integer from 1 to 8 and the R₄ radicals are each independently of the others as defined above;


$-\text{CX}=\text{CX}-$, wherein X is cyano, linear or branched C₁-C₈alkyl or di(linear or branched C₁-C₈alkyl)amino;

$-(\text{CH}_2)_q\text{-NR}_4\text{-(CH}_2\text{)}_q\text{-}$, wherein R₄ is as defined above and q is 1, 2, 3 or 4; or

a 1,2-cyclohexylene radical of formula:



or a 1,2-aryl radical of formula




wherein R₉ is hydrogen, SO₃H, CH₂OH or CH₂NH₂,

R, R₁ and R₁' are each independently of the others cyano; halogen; OR₄ or COOR₄ wherein R₄ is as defined above; nitro; linear or branched C₁-C₈alkyl; linear or branched partially fluorinated or perfluorinated C₁-C₈alkyl; or NHR₅, NR₅R₆ or N[®]R₅R₆R₇ wherein R₅, R₆ and R₇ are the same or different and are each hydrogen or linear or branched C₁-C₁₂alkyl or wherein R₅ and R₆ together with the nitrogen atom to which they are bonded form a 5-, 6- or 7-membered ring, which may contain further hetero atoms, or are linear or branched C₁-C₈alkyl-R₈ wherein R₈ is a radical OR₄, COOR₄ or NR₅R₆ as defined above or is NH₂ or N[®]R₅R₆R₇ wherein R₅, R₆ and R₇ are as defined above, R₂ and R₃ are each independently of the other hydrogen, linear or branched C₁-C₄alkyl, unsubstituted aryl or aryl that is substituted by cyano, by halogen, by OR₄ or COOR₄ wherein R₄ is hydrogen or linear or branched C₁-C₄alkyl, by nitro, by linear or branched C₁-C₈alkyl, by NHR₅ or NR₅R₆, wherein R₅ and R₆ are the same or different and are each linear or branched C₁-C₁₂alkyl or wherein R₅ and R₆ together with the nitrogen atom to which they are bonded form a 5-, 6- or 7-membered ring, which may contain further hetero atoms, by linear or branched C₁-C₈alkyl-R₇ wherein R₇ is an OR₄, COOR₄ or NR₅R₆ radical as defined above or is NH₂, or by N[®]R₅R₆R₇ wherein R₅, R₆ and R₇ are as defined above.

18. (new) Granules according to claim 17 that comprise as manganese complex a compound of formula (1) or (2) wherein Y is a radical of formula $-(\text{CH}_2)_r\text{-}$ wherein r is an integer from 1 to 4, or is a

radical of formula $-(R_4)_2-(CH_2)_p-C(R_4)_2-$ wherein p is a number from 0 to 3, and each R_4 , independently of the others, is hydrogen or C_1-C_4 alkyl, or is a 1,2-cyclohexylene radical or a 1,2-phenylene radical of formula:



19. **(new)** Granules according to claim 17 that comprise as manganese complex a compound of formula (1), (2) or (3) wherein the radicals R , R_1 and R_1' are hydrogen, OR_4 , $N(R_4)_2$ or $N^{\oplus}(R_4)_3$ and the R_4 groups in $N(R_4)_2$ or $N^{\oplus}(R_4)_3$ may be different and are each hydrogen or C_1-C_4 alkyl.

20. **(new)** Granules according to claim 17 that comprise as manganese complex a compound of formula (1), (2) or (3) wherein the radicals R_2 and R_3 are hydrogen, methyl, ethyl or unsubstituted phenyl.

21. **(new)** Granules according to claim 17 that comprise as manganese complex a compound of formula (1) or (2) wherein the anion A is a halide, perchlorate, sulfate, nitrate, hydroxide, BF_4^- , PF_6^- , carboxylate, acetate, tosylate or triflate.

22. **(new)** Granules according to claim 17 that comprise from 1 to 30 % by weight of manganese complex of formula (1), (2) or (3), based on the total weight of the granules.


23. **(new)** Granules according to claim 16 that comprise as dissolution restrainer an anionic dispersing agent, a non-ionic dispersing agent or a water-soluble organic polymer.

24. **(new)** Granules according to claim 23 that comprise as anionic dispersing agent a condensation product of a naphthalenesulfonic acid with formaldehyde, a sodium salt of a polymerised organic sulfonic acid, a (mono-/di-)alkylnaphthalenesulfonate, a polyalkylated polynuclear arylsulfonate, a sodium salt of a polymerised alkylbenzenesulfonic acid, a lignosulfonate, an oxylignosulfonate or a condensation product of naphthalenesulfonic acid with a polychloromethylidiphenyl.

25. **(new)** Granules according to claim 23 that comprise as non-ionic dispersing agent a compound selected from the group consisting of:

1. fatty alcohols having from 8 to 22 carbon atoms,
2. addition products of from 2 to 80 mol of alkylene oxide in which some alkylene oxide units are optionally replaced by substituted epoxides, with higher unsaturated or saturated monoalcohols, fatty acids, fatty amines or fatty amides having from 8 to 22 carbon atoms, or with benzyl alcohols, phenylphenols, benzylphenols or alkylphenols in which the alkyl radicals have at least 4 carbon atoms,
3. alkylene oxide condensation products,
4. ethylene oxide/propylene oxide adducts with diamines,
5. reaction products of a fatty acid having from 8 to 22 carbon atoms with a primary or secondary amine having at least one hydroxy-lower alkyl or lower alkoxy-lower alkyl group, or alkylene oxide addition products of such hydroxyalkyl-group-containing reaction products,
6. sorbitan esters or ethoxylated sorbitan esters,
7. addition products of propylene oxide with a tri- to hexa-hydric aliphatic alcohol having from 3 to 6 carbon atoms, and
8. fatty alcohol polyglycol mixed ethers.

26. **(new)** Granules according to claim 23 that comprise as non-ionic dispersing agent a surfactant of formula



wherein

R_{11} is C_8-C_{22} alkyl or C_8-C_{18} alkenyl;

R_{12} is hydrogen; C_1-C_4 alkyl; a cycloaliphatic radical having at least 6 carbon atoms or benzyl;

"alkylene" is an alkylene radical having from 2 to 4 carbon atoms and

n is a number from 1 to 60.

27. **(new)** Granules according claim 23 that comprise as water-soluble polymer a compound selected from the group consisting of:

polyethylene glycols, copolymers of ethylene oxide with propylene oxide, gelatin, polyacrylates, polymethacrylates, polyvinylpyrrolidones, vinylpyrrolidones, vinyl acetates, polyvinylimidazoles, polyvinylpyridine N-oxides, copolymers of vinylpyrrolidone with long-chained α -olefins, copolymers of vinylpyrrolidone with vinylimidazole, poly(vinylpyrrolidone/dimethylaminoethyl methacrylates), copolymers of vinylpyrrolidone/dimethylaminopropyl methacrylamides, copolymers of

vinylpyrrolidone/dimethylaminopropyl acrylamides, quaternised copolymers of vinylpyrrolidones and dimethylaminoethyl methacrylates, terpolymers of vinylcaprolactam/vinylpyrrolidone/-dimethylaminoethyl methacrylates, copolymers of vinylpyrrolidone and methacrylamidopropyl-trimethylammonium chloride, terpolymers of caprolactam/vinylpyrrolidone/dimethylaminoethyl methacrylates, copolymers of styrene and acrylic acid, polycarboxylic acids, polyacrylamides, carboxymethylcellulose, hydroxymethylcellulose, polyvinyl alcohols, optionally hydrolysed polyvinyl acetate, copolymers of ethyl acrylate with methacrylate and methacrylic acid, copolymers of maleic acid with unsaturated hydrocarbons and mixed polymerisation products of the said polymers.

28. **(new)** Granules according to claim 27 that comprise as organic polymer carboxymethylcellulose, a polyacrylamide, a polyvinyl alcohol, a polyvinylpyrrolidone, gelatin, a hydrolysed polyvinyl acetate, a copolymer of vinylpyrrolidone and vinyl acetate, a polyacrylate, a copolymer of ethyl acrylate with methacrylate and methacrylic acid or a polymethacrylate.

29. **(new)** Granules according to claim 16 that comprise the dissolution restrainer in an amount of from 10 to 95 % by weight based on the total weight of the granules.

30. **(new)** A washing agent formulation comprising

I) from 5 to 90 of an anionic surfactant A) and/or of a non-ionic surfactant B),

II) from 5 to 70 % of a builder substance C),

III) from 0.1 to 30 % of a peroxide D) and

IV) granules according to claim 17 in such an amount that the washing agent formulation comprises from 0.005 to 2 % of the pure manganese complex of formula (1), (2) or (3), the percentage figures in each case being percentages by weight based on the total weight of the washing agent.--

REMARKS

Claims 16-30 are the only claims pending. Said claims have been presented to reduce filing fees by eliminating multiple dependency, improper multiple dependency, improper multiple ranges for variables and to provide minor clarification.

The newly added claims are supported by originally filed claims 1-15. No new matter has been added.

Applicants aver that the claims are now in proper form for examination. An Action on the merits of the claims is respectfully awaited.

Respectfully submitted,



Kevin T. Mansfield
Agent for Applicants
Reg. No. 31,635

Ciba Specialty Chemicals Corporation
Patent Department
540 White Plains Road
P.O. Box 2005
Tarrytown, NY 10591-9005
(914) 785-7127
KTM22056PA

JAN 24 2002

Water-soluble granules of salen-type manganese complexes

The present invention relates to water-soluble granules of salen-type manganese complexes, to a process for the preparation thereof and to the use thereof as dye-transfer inhibitors in washing agent preparations.

A number of salen-type manganese complexes are already known to be suitable catalysts for oxidations with peroxy compounds, especially within the context of washing procedures. The use of certain manganese complexes as catalysts for preventing the redeposition of migrating dyes in peroxide-containing washing liquors is described in EP 902 083, but the action of those manganese complexes as dye-transfer inhibitors is not optimum under all washing conditions. A further problem is that the peroxy compound and/or the catalyst in the washing agent formulation decompose(s) during prolonged storage in a moist atmosphere.



Surprisingly, it has now been found that granules comprising a salen-type manganese complex and at least 10 % by weight of an anionic or non-ionic dissolution restrainer provide better inhibition of the redeposition of migrating dyes in washing liquors than is provided by the pure manganese complexes when the total amount of manganese complex entering into the washing liquor is the same in both cases. A further advantage of the granules is that the storage stability of peroxide-containing washing agent formulations comprising such granules is improved. In addition, these granules inhibit undesired colouration of the washing agent as a result of the gradual dissolution of the manganese complexes in one or more of the washing agent components.

The present invention accordingly relates to water-soluble granules of salen-type manganese complexes, comprising


- a) from 1 to 89 % by weight, preferably from 1 to 30 % by weight, of a water-soluble salen-type manganese complex,
- b) from 10 to 95 % by weight of a dissolution restrainer,
- c) from 0 to 20 % by weight of a further additive and
- d) from 1 to 15 % by weight of water, based on the total weight of the granules.

As manganese complexes for the granules according to the invention there come into consideration compounds that contain, complexed with manganese, from 1 to 3 saldimine groups, that is to say, groups obtainable by condensing unsubstituted or substituted salicylaldehydes with amines.

Especially suitable are compounds of formula



or



wherein

A is an anion;


m, n and p are each independently of the others 0, 1, 2 or 3,

R4 is hydrogen or linear or branched C1-C4alkyl,

- Y is a linear or branched alkylene radical of formula $-\text{C}(\text{R}_4)_2-$, wherein r is an integer from 1 to 8 and the R_4 radicals are each independently of the others as defined above;
- $-\text{CX}=\text{CX}-$, wherein X is cyano, linear or branched $\text{C}_1\text{-C}_8$ alkyl or di(linear or branched $\text{C}_1\text{-C}_8$ alkyl)amino;
- $-(\text{CH}_2)_q\text{-NR}_4\text{-(CH}_2)_q-$, wherein R_4 is as defined above and q is 1, 2, 3 or 4; or
- a 1,2-cyclohexylene radical of formula:



or a 1,2-aryl radical of formula



wherein R_9 is hydrogen, SO_3H , CH_2OH or CH_2NH_2 ,

R , R_1 and R_1' are each independently of the others cyano; halogen; OR_4 or COOR_4 wherein R_4 is as defined above; nitro; linear or branched $\text{C}_1\text{-C}_8$ alkyl; linear or branched partially fluorinated or perfluorinated $\text{C}_1\text{-C}_8$ alkyl; or NHR_5 , NR_5R_6 or $\text{N}^{\oplus}\text{R}_5\text{R}_6\text{R}_7$ wherein R_5 , R_6 and R_7 are the same or different and are each hydrogen or linear or branched $\text{C}_1\text{-C}_{12}$ alkyl or wherein R_5 and R_6 together with the nitrogen atom to which they are bonded form a 5-, 6- or 7-membered ring, which may contain further hetero atoms, or are linear or branched $\text{C}_1\text{-C}_8$ alkyl- R_8 wherein R_8 is a radical OR_4 , COOR_4 or NR_5R_6 as defined above or is NH_2 or $\text{N}^{\oplus}\text{R}_5\text{R}_6\text{R}_7$ wherein R_5 , R_6 and R_7 are as defined above,

R_2 and R_3 are each independently of the other hydrogen, linear or branched $\text{C}_1\text{-C}_4$ alkyl, unsubstituted aryl or aryl that is substituted by cyano, by halogen, by OR_4 or COOR_4 wherein R_4 is hydrogen or linear or branched $\text{C}_1\text{-C}_4$ alkyl, by nitro, by linear or branched $\text{C}_1\text{-C}_8$ alkyl, by NHR_5 or NR_5R_6 , wherein R_5 and R_6 are the same or different and are each linear or branched $\text{C}_1\text{-C}_{12}$ alkyl or wherein R_5 and R_6 together with the nitrogen atom to which they are bonded form a 5-, 6- or 7-membered ring, which may contain further hetero atoms, by linear or branched $\text{C}_1\text{-C}_8$ alkyl- R_7 wherein R_7 is an OR_4 , COOR_4 or NR_5R_6 radical as defined above or is NH_2 , or by $\text{N}^{\oplus}\text{R}_5\text{R}_6\text{R}_7$ wherein R_5 , R_6 and R_7 are as defined above.


When, in the compounds of formulae (1) and (3), R , R_1 , R_1' and/or R_8 are $\text{N}^{\oplus}\text{R}_5\text{R}_6\text{R}_7$ or R_2 and/or R_3 are $\text{N}^{\oplus}\text{R}_5\text{R}_6\text{R}_7$ -substituted aryl wherein R_5 , R_6 and R_7 are as defined above, the following anions are suitable for balancing the positive charge on the $\text{N}^{\oplus}\text{R}_5\text{R}_6\text{R}_7$ group:

halide, for example chloride, perchlorate, sulfate, nitrate, hydroxide, BF_4^- , PF_6^- , carboxylate, acetate, tosylate and triflate. Of those anions, bromide and chloride are preferred.

In compounds of formulae (1) and (3) in which n, m or p is 2 or 3, the radicals R, R₁ and R₁' have the same or different meanings.

When Y is a 1,2-cyclohexylene radical, it may be present in any of its stereoisomeric *cis/trans* forms.

Preferably, Y is a radical of formula $-(\text{CH}_2)_r-$ wherein r is an integer from 1 to 4, especially 2, or is a radical of formula $-\text{C}(\text{R}_4)_2-(\text{CH}_2)_p-\text{C}(\text{R}_4)_2-$ wherein p is a number from 0 to 3, especially 0, and each R₄, independently of the others, is hydrogen or C₁-C₄alkyl, especially hydrogen or methyl, or is a 1,2-cyclohexylene radical or a 1,2-phenylene radical of formula:



Halogen is preferably chlorine, bromine or fluorine, chlorine being especially preferred.

When n, m or p is 1, the groups R, R₁ and R₁' are preferably in the 4-position of the respective benzene ring except when R, R₁ or R₁' is nitro or COOR₄, in which case that group is preferably in the 5-position. When R, R₁ or R₁' is a N[⊕]R₅R₆R₇ group, that group is preferably in the 4- or 5-position.

When n, m or p is 2, the two R, R₁ or R₁' groups are preferably in the 4,6-position of the respective benzene ring except when they are nitro or COOR₅, in which case the two groups are preferably in the 3,5-position.

When R, R₁ or R₁' is di(C₁-C₁₂alkyl)amino, the alkyl group may be straight-chain or branched. Preferably, it contains from 1 to 8, especially from 1 to 3, carbon atoms.

Preferably, the radicals R , R_1 and R_1' are hydrogen, OR_4 , $N(R_4)_2$ or $N^+(R_4)_3$, wherein the R_4 groups in $N(R_4)_2$ or $N^+(R_4)_3$ may be different and are hydrogen or C_1 - C_4 alkyl, especially methyl, ethyl or isopropyl.

The radicals R_2 and R_3 are especially hydrogen, methyl, ethyl or unsubstituted phenyl.

Aryl is, for example, naphthyl or, especially, phenyl.

When R_5 and R_6 together with the nitrogen atom to which they are bonded form a 5-, 6- or 7-membered ring, the ring is especially a pyrrolidine, piperidine, morpholine or piperazine ring. The piperazine ring may be substituted, for example by alkyl, at the nitrogen atom that is not bonded to the phenyl or alkyl radical.

Suitable anions A include, for example, halide, such as chloride or bromide, perchlorate, sulfate, nitrate, hydroxide, BF_4^- , PF_6^- , carboxylate, acetate, tosylate and triflate. Of those anions, chloride, bromide and acetate are preferred.

The compounds of formulae (1), (2) and (3) are known or can be prepared in a manner known *per se*. The manganese complexes are prepared from the corresponding ligands and a manganese compound. Such preparation procedures are described, for example, in US Patents 5 281 578 and 4 066 459 and by Bernardo *et al.*, Inorg. Chem. 45 (1996) 387.

Preferred formulations of the granules comprise from 1 to 90 % by weight, especially from 1 to 30 % by weight, of salen-type manganese complex of formula (1), (2) or (3), based on the total weight of the granules.

Instead of a single, homogeneous manganese complex of formula (1), (2) or (3) it is also possible to use mixtures of two or more manganese complexes of formula (1), (2) or (3). Mixtures of one or more manganese complexes of formula (1), (2) or (3) and one or more salen-type ligands can also be used. Salen-type ligands suitable for such mixtures include all ligands that are used as starting compounds in the preparation of the manganese complexes of formula (1), (2) and (3).

As dissolution restrainers for the granules according to the invention there come into consideration compounds that cause the manganese complexes to dissolve in water more slowly than they would without the dissolution restrainers. The following, for example, come into consideration:

1. anionic dispersing agents,
2. non-ionic dispersing agents and
3. water-soluble organic polymers.

The anionic dispersing agents used are, for example, the commercially available water-soluble anionic dispersing agents for dyes, pigments etc.. The following products, especially, come into consideration: condensation products of aromatic sulfonic acids and formaldehyde, condensation products of aromatic sulfonic acids with unsubstituted or chlorinated diphenylene or diphenyl oxides and, optionally, formaldehyde, (mono-/di-)alkylnaphthalene-sulfonates, sodium salts of polymerised organic sulfonic acids, sodium salts of polymerised alkylnaphthalenesulfonic acid, sodium salts of polymerised alkylbenzenesulfonic acid, alkylarylsulfonates, sodium salts of alkyl polyglycol ether sulfates, polyalkylated polynuclear arylsulfonates, methylene-linked condensation products of arylsulfonic acids and hydroxy-aryl sulfonic acids, sodium salts of dialkylsulfosuccinic acid, sodium salts of alkyl diglycol ether sulfates, sodium salts of polynaphthalenemethanesulfonates, ligno- or oxygigno-sulfonates and heterocyclic polysulfonic acids.

The following anionic dispersing agents are especially suitable: condensation products of naphthalenesulfonic acids with formaldehyde, sodium salts of polymerised organic sulfonic acids, (mono-/di-)alkylnaphthalenesulfonates, polyalkylated polynuclear arylsulfonates, sodium salts of polymerised alkylbenzenesulfonic acid, lignosulfonates, oxygignosulfonates and condensation products of naphthalenesulfonic acid with a polychloromethyldiphenyl.


Suitable non-ionic dispersing agents are especially compounds having a melting point of at least 35°C that are emulsifiable, dispersible or soluble in water. They include, for example, the following compounds:

1. fatty alcohols having from 8 to 22 carbon atoms, especially cetyl alcohol,
2. addition products of preferably from 2 to 80 mol of alkylene oxide, especially ethylene oxide, in which individual ethylene oxide units may have been replaced by substituted epoxides, such as styrene oxide and/or propylene oxide, with higher unsaturated or

saturated monoalcohols, fatty acids, fatty amines or fatty amides having from 8 to 22 carbon atoms, or with benzyl alcohols, phenylphenols, benzylphenols or alkylphenols in which the alkyl radicals have at least 4 carbon atoms,

3. alkylene oxide condensation products, especially propylene oxide condensation products (block polymers),
4. ethylene oxide/propylene oxide adducts with diamines, especially ethylenediamine,
5. reaction products of a fatty acid having from 8 to 22 carbon atoms with a primary or secondary amine having at least one hydroxy-lower alkyl or lower alkoxy-lower alkyl group, or alkylene oxide addition products of such hydroxyalkyl-group-containing reaction products,
6. sorbitan esters, preferably having long-chained ester groups, or ethoxylated sorbitan esters, such as, for example, polyoxyethylene-sorbitan monolaurate having from 4 to 10 ethylene oxide units or polyoxyethylene-sorbitan trioleate having from 4 to 20 ethylene oxide units,
7. addition products of propylene oxide with a tri- to hexa-hydric aliphatic alcohol having from 3 to 6 carbon atoms, for example glycerol or pentaerythritol, and
8. fatty alcohol polyglycol mixed ethers, especially addition products of from 3 to 30 mol of ethylene oxide and from 3 to 30 mol of propylene oxide with aliphatic monoalcohols having from 8 to 22 carbon atoms.

Non-ionic dispersing agents that are especially suitable are surfactants of formula



wherein

R_{11} is C_8-C_{22} alkyl or C_8-C_{18} alkenyl;

R_{12} is hydrogen; C_1-C_4 alkyl; a cycloaliphatic radical having at least 6 carbon atoms or benzyl;

"alkylene" is an alkylene radical having from 2 to 4 carbon atoms and

n is a number from 1 to 60.


The substituents R_{11} and R_{12} in formula (4) are advantageously the hydrocarbon radical of an unsaturated or, preferably, saturated aliphatic monoalcohol having from 8 to 22 carbon atoms. The hydrocarbon radical may be straight-chain or branched. Preferably, R_{11} and R_{12} are each independently of the other an alkyl radical having from 9 to 14 carbon atoms.

As saturated aliphatic monoalcohols there come into consideration natural alcohols, such as, for example, lauryl alcohol, myristyl alcohol, cetyl alcohol and stearyl alcohol, as well as synthetic alcohols, such as, for example, 2-ethylhexanol, 1,1,3,3-tetramethylbutanol, octan-2-ol, isononyl alcohol, trimethylhexanol, trimethylnonyl alcohol, decanol, C₉-C₁₁oxoalcohol, tridecyl alcohol, isotridecyl alcohol and linear primary alcohols (Alfols) having from 8 to 22 carbon atoms. Some examples of such Alfols are Alfol (8-10), Alfol (9-11), Alfol (10-14), Alfol (12-13) and Alfol (16-18). ("Alfol" is a registered trade mark).

Unsaturated aliphatic monoalcohols are, for example, dodecenyl alcohol, hexadecenyl alcohol and oleyl alcohol.


The alcohol radicals may be used individually or in the form of mixtures of two or more components, such as, for example, mixtures of alkyl and/or alkenyl groups derived from soybean fatty acids, palm-kernel fatty acids or tallow oils.

(Alkylene-O) chains are preferably divalent radicals of formula



Examples of a cycloaliphatic radical are cycloheptyl, cyclooctyl and, preferably, cyclohexyl.

As non-ionic dispersing agents there preferably come into consideration surfactants of formula



wherein

R₁₃ is C₈-C₂₂alkyl;


R₁₄ is hydrogen or C₁-C₄alkyl;

Y₁, Y₂, Y₃ and Y₄ are each independently of the others hydrogen, methyl or ethyl;

n₂ is a number from 0 to 8; and

n₃ is a number from 2 to 40.

Further important non-ionic dispersing agents correspond to the formula



wherein

R_{15} is C_8 - C_{14} alkyl;

R_{16} is C_1 - C_4 alkyl;

Y_5, Y_6, Y_7 and Y_8 are each independently of the others hydrogen, methyl or ethyl, one of the radicals Y_5, Y_6 and one of the radicals Y_7, Y_8 always being hydrogen; and n_4 and n_5 are each independently of the other an integer from 4 to 8.

The non-ionic dispersing agents of formulae (4) to (6) can be used in the form of mixtures. There come into consideration as surfactant mixtures, for example, non-end-group-terminated fatty alcohol ethoxylates of formula (4), that is to say, compounds of formula (4) wherein

R_{11} is C_8 - C_{22} alkyl,

R_{12} is hydrogen and

the alkylene-O chain is the radical $-(CH_2-CH_2-O)-$

as well as end-group-terminated fatty alcohol ethoxylates of formula (6).

As examples of non-ionic dispersing agents of formulae (4), (5) and (6) there may be mentioned reaction products of a C_{10} - C_{13} fatty alcohol, for example a C_{13} oxoalcohol, with from 3 to 10 mol of ethylene oxide, propylene oxide and/or butylene oxide, or the reaction product of 1 mol of a C_{13} fatty alcohol with 6 mol of ethylene oxide and 1 mol of butylene oxide, it being possible for the addition products in each case to be terminated by a C_1 - C_4 alkyl end group, preferably methyl or butyl.

The dispersing agents may be used individually or in the form of mixtures of two or more dispersing agents.

Instead of or in addition to the anionic or non-ionic dispersing agent, the granules according to the invention may comprise a water-soluble organic polymer as dissolution restrainer.

Such polymers may be used individually or in the form of mixtures of two or more polymers. Preferably, such a polymer is added for the purpose of improving the mechanical stability of the granules and/or when, during later use of the granules in the washing agent, the dissolution of the salen-type manganese complex in the washing liquor is to be controlled, and/or when an enhanced action as dye inhibitor is desired.

As water-soluble polymers there come into consideration, for example, polyethylene glycols, copolymers of ethylene oxide with propylene oxide, gelatin, polyacrylates, polymethacrylates, polyvinylpyrrolidones, vinylpyrrolidones, vinyl acetates, polyvinylimidazoles, polyvinylpyridine N-oxides, copolymers of vinylpyrrolidone with long-chained α -olefins, copolymers of vinylpyrrolidone with vinylimidazole, poly(vinylpyrrolidone/dimethylaminoethyl methacrylates), copolymers of vinylpyrrolidone/dimethylaminopropyl methacrylamides, copolymers of vinylpyrrolidone/dimethylaminopropyl acrylamides, quaternised copolymers of vinylpyrrolidones and dimethylaminoethyl methacrylates, terpolymers of vinylcaprolactam/vinylpyrrolidone/dimethylaminoethyl methacrylates, copolymers of vinylpyrrolidone and methacrylamidopropyl-trimethylammonium chloride, terpolymers of caprolactam/vinylpyrrolidone/dimethylaminoethyl methacrylates, copolymers of styrene and acrylic acid, polycarboxylic acids, polyacrylamides, carboxymethylcellulose, hydroxymethylcellulose, polyvinyl alcohols, optionally hydrolysed polyvinyl acetate, copolymers of ethyl acrylate with methacrylate and methacrylic acid, copolymers of maleic acid with unsaturated hydrocarbons and mixed polymerisation products of the said polymers.

Among those organic polymers, special preference is given to carboxymethylcellulose, polyacrylamides, polyvinyl alcohols, polyvinylpyrrolidones, gelatin, hydrolysed polyvinyl acetate, copolymers of vinylpyrrolidone and vinyl acetate and also polyacrylates, copolymers of ethyl acrylate with methacrylate and methacrylic acid and polymethacrylates.

The dissolution restrainers are used in an amount of from 10 to 95 % by weight, preferably from 15 to 85 % by weight and especially from 25 to 75 % by weight, based on the total weight of the granules.

The granules according to the invention may comprise further additives, for example wetting agents, water-insoluble or water-soluble dyes or pigments and also fillers and optical brighteners. Such additives are present in an amount of from 0 to 20 % by weight, based on the total weight of the granules.

The granules according to the invention are prepared, for example, starting from:

- a) a solution or suspension with a subsequent drying/forming step or
- b) a suspension of the active ingredient in a melt, with subsequent forming and solidification.

a) First, the anionic or non-ionic dispersing agent and/or the polymer and, as appropriate, the further additives are dissolved in water and stirred, optionally with heating, until a homogeneous solution is obtained. The salen-type manganese complex is then dissolved or suspended in the resulting aqueous solution. The solids content of the solution should preferably be at least 30 % by weight, more especially from 40 to 50 % by weight, based on the total weight of the solution. The viscosity of the solution is preferably less than 200 mPas.

In a drying step all the water, with the exception of a residual amount, is then removed from the so-prepared aqueous solution comprising the salen-type manganese complex, solid particles (granules) simultaneously being formed. Known methods are suitable for producing the granules from the aqueous solution. In principle, both methods with continuous operation and those with discontinuous operation are suitable. Preference is given to continuous processes, especially spray-drying granulation methods and fluidised-bed granulation methods.

Spray-drying methods in which the active ingredient solution is sprayed into a chamber in which hot air is being circulated are especially suitable. The atomisation of the solution is carried out, for example, using unitary or binary nozzles or is brought about by the spinning effect of a rapidly rotating disc. In order to increase the particle size, the spray-drying procedure can be combined with an additional agglomeration of the liquid particles with solid nuclei in a fluidised bed integrated in the chamber (so-called fluid-spray). The fine particles (<100 µm) obtained by a conventional spray-drying method may, if necessary after being separated from the exhaust air flow, be fed directly, without being further treated, to the atomizing cone of the spray-dryer atomizer, as nuclei for the purpose of agglomeration with the liquid droplets of the active ingredient.

During the granulation step, the water can rapidly be removed from the solutions comprising the salen-type manganese complex, dissolution restrainer and further additives, and it is expressly intended that agglomeration of the droplets forming in the atomizing cone, or the agglomeration of droplets with solid particles, will take place.

If necessary, the granules formed in the spray-dryer are separated off in a continuous process, for example by means of a sieving operation. The fine particles and the oversize particles are either recycled in the process directly (without being dissolved) or are dissolved in the liquid active ingredient formulation and then granulated again.

The granules according to the invention are resistant to abrasion, low in dust, are free-flowing and easily metered. A distinguishing feature is that their rate of dissolution in water is controllable by the composition of the formulation. They are used especially in washing agent formulations as dye-transfer inhibitors. They can be added directly to a washing agent formulation at the desired concentration of the salen-type manganese complex. The present invention relates also to that use.

Where the coloured appearance of the granules in the washing agent is to be suppressed, that can be achieved, for example, by embedding the granules in droplets consisting of a whitish meltable substance ("water-soluble wax"), or by adding a white pigment (e.g. TiO_2) to the granule formulation or, preferably, by encasing the granules with a melt consisting, for example, of a water-soluble wax, as described in EP-B-0 323 407 B1, a white solid (e.g. titanium dioxide) being added to the melt in order to reinforce the masking effect of the casing.

b) Prior to granulation of the melt, the salen-type manganese complex is dried in a separate step and, if necessary, dry-ground in a mill so that all solid particles are $< 50 \mu\text{m}$. The drying is carried out in an apparatus customary for that purpose, for example in a paddle dryer, a vacuum cabinet or a freeze-dryer.

The finely particulate manganese complex is suspended in the molten carrier material and the suspension is homogenised. The desired granules are prepared from the suspension in a forming step with simultaneous solidification of the melt. The selection of a suitable melt-granulation method is dependent upon the desired size of the granules. In principle, any method that allows the production of granules of a particle size of from 0.1 to 4 mm is suitable. Such methods include droplet-dispensing processes (with solidification on a cooling belt), prilling (gas/liquid cooling medium) and flake formation with a subsequent comminution step, the granulating apparatus being operated continuously or discontinuously.

Where the coloured appearance of the granules in the washing agent is to be suppressed, there can also be suspended in the melt, in addition to the manganese complex, white or coloured pigments (e.g. titanium dioxide) that impart the desired colour appearance to the granules after solidification.

The present invention accordingly relates also to washing agent formulations comprising I) from 5 to 90 %, preferably from 5 to 70 %, A) of an anionic surfactant and/or B) of a non-ionic surfactant, II) from 5 to 70 %, preferably from 5 to 50 %, especially from 5 to 40 %, C) of a builder substance, III) from 0.1 to 30 %, preferably from 1 to 12 %, D) of a peroxide and IV) E) granules according to the invention in such an amount that the washing agent formulation comprises from 0.005 to 2 %, preferably from 0.02 to 1 %, especially from 0.1 to 0.5 %, of the pure manganese complex of formula (1), (2) or (3). In each case, the percentage figures are percentages by weight, based on the total weight of the washing agent.

The washing agent may be in solid or liquid form, but in liquid form it is preferably a non-aqueous washing agent containing not more than 5 % by weight, preferably from 0 to 1 % by weight, of water and comprising as base a suspension of a builder substance in a non-ionic surfactant, for example as described in GB-A-2 158 454.

The washing agent is preferably, however, in the form of a powder or granules.

The powder or granules can be produced, for example, by first of all preparing a starting powder by spray-drying an aqueous suspension comprising all of the components listed above, with the exception of components D) and E), and then adding the dry components D) and E) and mixing everything together.

It is also possible to start with an aqueous suspension that comprises components A) and C) but not component B) or only a proportion of component B). The suspension is spray-dried and then component E) is mixed with component B) and the mixture is added to the suspension, and subsequently component D) is admixed dry.

10043045-012402
204220-2404001

Preferably, the components are mixed together in such amounts that a solid compact washing agent in the form of granules is obtained that has a specific weight of at least 500 g/l.

In a further preferred embodiment, the washing agent is prepared in three steps. In the first step a mixture of anionic surfactant (and, if desired, a small amount of non-ionic surfactant) and builder substance is prepared. In the second step that mixture is sprayed with the bulk of the non-ionic surfactant, and then in the third step peroxide, catalyst as appropriate, and the granules according to the invention are added. That method is normally carried out in a fluidised bed.

In a further preferred embodiment, the individual steps are not carried out completely separately, resulting in a certain amount of overlap between them. Such a method is usually carried out in an extruder, in order to obtain granules in the form of "megapearls".

The anionic surfactant A) may be, for example, a sulfate, sulfonate or carboxylate surfactant or a mixture of such surfactants.

Preferred sulfates are those having from 12 to 22 carbon atoms in the alkyl radical, where appropriate in combination with alkyl ethoxysulfates in which the alkyl radical contains from 10 to 20 carbon atoms.

Preferred sulfonates include, for example, alkylbenzenesulfonates having from 9 to 15 carbon atoms in the alkyl radical and/or alkylnaphthalenesulfonates having from 6 to 16 carbon atoms in the alkyl radical.

The cation in the anionic surfactants is preferably an alkali metal cation, especially sodium.

Preferred carboxylates are alkali metal sarcosinates of formula $R-CO-N(R^1)-CH_2COOM^1$, wherein R is alkyl or alkenyl having from 8 to 18 carbon atoms in the alkyl or alkenyl radical, R^1 is C_1 - C_4 alkyl and M^1 is an alkali metal.

The non-ionic surfactant B) may be, for example, a condensation product of from 3 to 8 mol of ethylene oxide with 1 mol of primary alcohol that contains from 9 to 15 carbon atoms.

There come into consideration as builder substance C), for example, alkali metal phosphates, especially tripolyphosphates, carbonates or bicarbonates, especially the sodium salts thereof, silicates, aluminium silicates, polycarboxylates, polycarboxylic acids, organic phosphonates, aminoalkylenepoly(alkylenephosphonates) and mixtures of such compounds.

Especially suitable silicates are sodium salts of crystalline layer silicates of the formula $\text{NaHSi}_t\text{O}_{2t+1} \cdot p\text{H}_2\text{O}$ or $\text{Na}_2\text{Si}_t\text{O}_{2t+1} \cdot p\text{H}_2\text{O}$, wherein t is a number from 1.9 to 4 and p is a number from 0 to 20.

Among the aluminium silicates, preference is given to those obtainable commercially under the names zeolite A, B, X and HS and also to mixtures of two or more of those components.

Among the polycarboxylates, preference is given to polyhydroxycarboxylates, especially citrates, and acrylates and also copolymers thereof with maleic anhydride.

Preferred polycarboxylic acids are nitrilotriacetic acid, ethylenediaminetetraacetic acid and ethylenediamine disuccinate either in racemic form or in the enantiomerically pure S,S form.

Especially suitable phosphonates and aminoalkylenepoly(alkylenephosphonates) include alkali metal salts of 1-hydroxyethane-1,1-diphosphonic acid, nitrilotris(methylenephosphonic acid), ethylenediaminetetramethylenephosphonic acid and diethylenetriaminepentamethylenephosphonic acid.

As the peroxide component D) there come into consideration, for example, the organic and inorganic peroxides known in the literature and available commercially that bleach textiles at conventional washing temperatures, for example at from 10 to 95°C.

The organic peroxides are, for example, mono- or poly-peroxides, especially organic peracids or salts thereof, such as phthalimidoperoxycaproic acid, peroxybenzoic acid, diperoxydodecanedioic acid, diperoxynonanedioic acid, diperoxydecanedioic acid, diperoxyphthalic acid or salts thereof.

Preference is given, however, to the use of inorganic peroxides, such as, for example, persulfates, perborates, percarbonates and/or persilicates. It will be understood that it is also possible to use mixtures of inorganic and/or organic peroxides. The peroxides may be in a variety of crystalline forms and may have different water contents, and they may also be used together with other inorganic or organic compounds in order to improve their storage stability.

The peroxides are added to the washing agent preferably by mixing the components together, for example using a screw metering system and/or a fluidised bed mixer.

The washing agent may comprise, in addition to the granules according to the invention, one or more optical brighteners, for example from the group bistriazinylaminostilbene-disulfonic acid, bistriazolylstilbenedisulfonic acid, bisstyrylbiphenyl or bisbenzofuranyl-biphenyl, a bisbenzoxalyl derivative, bisbenzimidazolyl derivative, coumarin derivative or a pyrazoline derivative.


The washing agents may furthermore comprise suspending agents for dirt, e.g. sodium carboxymethylcellulose, pH regulators, e.g. alkali metal or alkaline earth metal silicates, foam regulators, e.g. soap, salts for regulating the spray-drying and the granulating properties, e.g. sodium sulfate, perfumes and, optionally, antistatic agents and softeners, enzymes, such as amylase, bleaching agents, pigments and/or toning agents. It will be understood that such components must be stable towards the bleaching agent used.

Further preferred additives for the washing agents according to the invention are polymers that, during the washing of textiles, inhibit staining caused by dyes in the washing liquor that have been released from the textiles under the washing conditions. Such polymers are preferably polyvinylpyrrolidones, polyvinylimidazoles or polyvinylpyridine N-oxides which may have been modified by the incorporation of anionic or cationic substituents, especially those having a molecular weight in the range from 5000 to 60 000, more especially from


10 000 to 50 000. Such polymers are used preferably in an amount of from 0.05 to 5 % by weight, especially from 0.2 to 1.7 % by weight, based on the total weight of the washing agent.

In addition, the washing agents according to the invention may also comprise so-called perborate activators, such as, for example, TAED, SNOBS or TAGU. Preference is given to TAED, which is preferably used in an amount of from 0.05 to 5 % by weight, especially from 0.2 to 1.7 % by weight, based on the total weight of the washing agent.


The following Examples serve to illustrate the invention without the invention being limited thereto. Parts and percentages are by weight unless specified otherwise. The manganese complexes used in the Examples are the compounds of formulae (1a), (1b) and (3a):



(1a)



(1b)



5 different granules are used in the Examples.

Example 1: 150 g of polyvinyl alcohol (PVA) (MW=15 000) are dissolved in 850 g of water at approximately 50°C. After the PVA has dissolved completely, 7.5 g of filter cake (active content = 45.3 %) of the manganese complex of structure (1a) are added to the solution and the complex is dissolved with stirring.

The solution is then spray-dried in a spray-dryer equipped with a binary nozzle. The exhaust-air temperature is 120°C at a feed-air temperature of 210°C. Free-flowing granules having a mean particle size of 15 µm and a residual water content of 10 % are obtained. The granules produced in that manner contain 2 % of the manganese complex of structure (1a).

Examples 2 to 4:

Granules having the following compositions are produced according to the same procedure:

Ex. No.	Manganese complex	% by wt. of manganese complex	Polymer	% by wt. of polymer	% by wt. of residual moisture in the granules
2	structure (1a)	2	sodium carboxy-methylcellulose	87	11
3	structure (1a)	2	gelatin	84	14
4	structure (1a)	2	copolymer of ethyl acrylate with methacrylate and methacrylic acid	91	7

Example 5:

Moist filter cake of the manganese complex of structure (1a) is dried in a vacuum cabinet to a residual moisture content of 5.2 %. The dried manganese complex is ground in a laboratory mill to a mean particle size of 36 μm .

200 g of polyethylene glycol 8000 (melting point 63°C) are introduced as initial charge into a double-walled vessel equipped with a stirrer and a heatable outlet (modified to form a nozzle having a diameter of 0.8 mm). The polyethylene glycol is heated to 120°C under nitrogen. 2.042 g of the ground manganese complex of structure (1a) are stirred into the hot melt and the suspension is homogenised for a further 30 minutes.

The hot suspension is slowly dispensed in droplets onto a cooled rotating metal plate. The hot droplets solidify in approximately 10 seconds to form the desired granules having an average diameter of 2 mm. The size of the granules can be controlled, for example, by the temperature of the melt. The granules contain 2 % of the manganese complex of structure (1a).

Example 6: Release of the manganese complex into solution:

The rate at which the granules release the manganese complex into an alkaline solution at 40°C is determined as follows:

0.1107 g of granules is added with stirring, at time 0, to 100 ml of borax buffer (pH=10, 0.03 g/l of disodium tetraborate and 0.042 g/l of sodium hydroxide). After set intervals, a

sample of the solution is taken and an absorption spectrum thereof is measured. The manganese complex has an absorption band at 405 nm. The optical density of a solution of 0.022 g/l of the fully dissolved catalyst at 405 nm is 1.6. The Table below shows the results. It will be seen from the Table that the various granules release the manganese complex into the solution in a slow and controlled manner.

Granules from Example	Optical density at 405 nm							
	1 min	7 min	13 min	19 min	25 min	31 min	60 min	120 min
1	0.06	0.34	0.60	0.77	0.92	1.02	1.26	1.35
2	-	0.24	0.40	0.55	0.68	0.78	1.05	1.23
3	0.49	1.41	1.57	1.57	1.57	1.57	1.57	-
4	0.86	1.11	1.12	1.14	1.15	1.16	1.20	-
5	0.77	1.63	1.63	1.63	1.63	1.63	1.63	-

Example 7: In order to examine the effectiveness of the granules as dye-transfer inhibitors, the DTI activity is determined. The DTI (dye transfer inhibition) activity a is defined as the following percentage:

$$a = ([Y(E) - Y(A)] / [Y(W) - Y(A)]) * 100$$

wherein $Y(W)$, $Y(A)$ and $Y(E)$ denote the CIE brightness values of the white material, of the material treated without the addition of dye-transfer inhibitor and of the material treated with the addition of dye-transfer inhibitor, respectively. $a=0$ denotes a completely ineffective product, which when added to the washing liquor allows the dye transfer to proceed freely, whereas $a=100\%$ corresponds to a perfect dye-transfer inhibitor, which completely prevents staining of the white material.

The following test system is used to obtain the test data: 5 g of white cotton fabric are treated in 80 ml of washing liquor. The liquor comprises the standard washing agent ECE phosphate-free (456 IEC) EMPA, Switzerland, in a concentration of 7.5 g/l, 8.6 mmol/l of H_2O_2 and 5 g of cotton fabric dyed with dyestuff R Bk 5 (Reactive Black 5). The washing procedure is carried out in a beaker in a LINITEST apparatus for 30 min. at 40°C. The dye-transfer inhibitor is added in the amount indicated in each case. The reflection spectra of the

specimens are measured using a SPECTRAFLASH 2000 and converted into brightness values (D65/10) by standard CIE procedure.

It will be seen from the Table below that the granules exhibit a significantly better DTI activity than the pure manganese complex, although the absolute amount of pure manganese complex metered in is the same in all 6 experiments.

Dye-transfer inhibitor	0.00177 g (1a) *	0.0886 g of granules from Example 1	0.0886 g of granules from Example 2	0.0886 g of granules from Example 3	0.0886 g of granules from Example 4	0.0886 g of granules from Example 5
a (%)	0	38	57	27	18	58

* The indicated amount of pure manganese complex is metered in in the form of a concentrated methanolic solution. See Example 1 for the definition of dye-transfer inhibitors.

Example 8: The following test system is used to obtain the test data: 7.5 g of white cotton fabric are treated for 30 min. at 40°C in 80 ml of washing liquor. The liquor comprises the standard washing agent ECE phosphate-free (456 IEC) EMPA, Switzerland, in a concentration of 7.5 g/l and 8.6 mmol/l of H₂O₂. R Bk 5 in the 133 % formulation is used as dyestuff. Using a computer-controlled feed pump, the dyestuff is slowly metered in during the washing procedure in the form of a concentrated solution. In that way, the slow bleeding of dyes from coloured textiles is simulated. The concentration of dyestuff in the washing liquor as a function of time (K(t), K in mg/l, t in min.), without dye-transfer inhibitor and without fabric, is described by the function:

$$K(t) = 4.9 \cdot (1 - \exp(-0.059 \cdot t)) + 8.0 \cdot (1 - \exp(-1.46 \cdot t))$$

The concentration of dyestuff after 30 min. is accordingly 12 mg/l. The dye-transfer inhibitor is added at the beginning of the experiment in the amount indicated in each case. The reflection spectra of the specimens are measured using a SPECTRAFLASH 2000 and converted into brightness values (D65/10) by standard CIE procedure.

It will be seen from the Table below that the granules exhibit a significantly better DTI activity [see Example 7 for the definition of *a* (%)] than the pure manganese complex, although the absolute amount of pure manganese complex metered in is the same in all 6 experiments.

Dye-transfer inhibitor	0.00177 g of (1a) *	0.0886 g of granules from Example 1	0.0886 g of granules from Example 2	0.0886 g of granules from Example 3	0.0886 g of granules from Example 4	0.0886 g of granules from Example 5
<i>a</i> (%)	24	56	72	48	56	72

* The indicated amount of pure manganese complex was metered in in the form of a concentrated methanolic solution. See Example 1 for the definition of dye-transfer inhibitors.

Example 9: 0.1 g of compound (1b), 0.25 g of the dispersing agent 1618 (see below) and 4.65 g of the polymer PEG 8000 (see below) are melted at 80°C and the melt is stirred until homogeneous. Using a plastics pipette, small amounts of the melt are dispensed in droplets onto a cooled metal plate. The solidified droplets have an average size of approximately 5 mm.

Examples 10 to 30: The following formulations (see Table below) are prepared as described in Example 9. The compositions of the solid formulations are given in percent by weight.

2041210*54084001

Example	(1a)	(1b)	Disp. 1618	Pluronic F-108	Lutensol AT 25	Lutensol AT 50	PEG 8000	PEG 20 000
10		2					98	
11		2	10				88	
12		2	15				83	
13		2		5			93	
14		2			5		93	
15		10					90	
16		2	98					
17		2		98				
18		2			98			
19	2		98					
20	2			98				
21	2				98			
22	2		5				93	
23	2			5			93	
24	2				5		93	
25	2						20	78
26	2						78	20
27		2					20	78
28		2					78	20
29		10				90		
30	10					90		

--Dispersing agent 1618 = Marlipal 1618 = $\text{RO}(\text{CH}_2\text{CH}_2\text{O})_{25}\text{H}$, R = saturated linear $\text{C}_{16}\text{C}_{18}$ fatty alcohol (Hüls)

--Pluronic F-108 = EO/PO block polymer, M = 15 500 (BASF)

--Lutensol AT 25 = $\text{RO}(\text{CH}_2\text{CH}_2\text{O})_{25}\text{H}$, R = saturated linear $\text{C}_{16}\text{C}_{18}$ fatty alcohol (BASF)

--Lutensol AT 50 = $\text{RO}(\text{CH}_2\text{CH}_2\text{O})_{50}\text{H}$, R = saturated linear $\text{C}_{16}\text{C}_{18}$ fatty alcohol (BASF)

--PEG 20 000 = polyethylene glycol, Mr = approx. 16 000 - 24 000 (Fluka)

Example 31: A homogeneous suspension of 0.3 g of compound (1b) and 14.7 g of the dissolution restrainer Klucel E (see below) in 135 ml of deionised water is obtained after stirring for from 15 to 30 minutes. The suspension is dried at 80°C and 120 mbar for three days. The formulation is cooled to -73°C and pulverised in a mortar.

Examples 32 to 41: The following formulations (see Table below) are prepared as described in Example 31. The compositions of the solid formulations are given in percent by weight.

Example	(1a)	(1b)	Klucel E	PVP K-30	Acrysol A-3	Acrysol A-5	Glascol E-11
32		2	98				
33		2		98			
34	2		98				
35	2			98			
36	10				90		
37	10					90	
38	10						90
39		10			90		
40		10				90	
41		10					90

—Klucel E = hydroxypropyl cellulose, MW = 80 000 (Aqualon Company)

—PVP K-30 = polyvinylpyrrolidone, Mr = 80 000 (Erne Chemie)

—Acrysol A-3 = polyacrylic acid, MM < 150 000 (Rohm and Haas)

—Acrysol A-5 = polyacrylic acid, MM < 300 000 (Rohm and Haas)

—Glascol E-11 = polyacrylic acid, MM approx. 250 000 (Ciba Spezialitätenchemie)

Example 42: A homogeneous suspension comprising 1 g of compound (1a) and 9 g of the dissolution restrainer PVP K-30 in 115 g of deionised water is prepared by stirring for from 15 to 30 minutes. While being continuously rotated in a methylene chloride/dry-ice bath

(approximately -73°C), the suspension is frozen in the form of a thin layer in a 1-litre pear-shaped flask and is then lyophilised.

Examples 43 to 52: The following formulations (see Table below) are prepared as described in Example 42. The compositions of the solid formulations are given in percent by weight.

Example	(1a)	(1b)	PVP K-30	Acrysol A-1	Acrysol A-3	Acrysol A-5	Good- rite K-702
43		2	98				
44	2		98				
45		10		90			
46		10			90		
47		10				90	
48		10					90
49	10			90			
50	10				90		
51	10					90	
52	10						90

--Acrysol A-1 = polyacrylic acid, MM < 50 000 (Rohm and Haas)

--Good-rite K-702 = polyacrylic acid, Mw = 243 000 (BFGoodrich)

Example 53: A formulation is prepared as described in Example 9 from 10 % by weight of compound (3a) and 90 % by weight of Lutensol AT 50.

Examples 54 - 60: The test data are obtained in a manner analogous to that described in Example 8 of the Application. In all of the Examples below, the concentration of pure catalyst (1a) in the washing liquor is 50 μ M (= 22 mg/l).



Example	Formulation from Example	DTI effect a (%)
54	10	61
55	30	62

56	34	73
57	35	86
58	44	73
59	51	65
60	52	59


10048045-012402

What is claimed is:

1. Water-soluble granules of salen-type manganese complexes, comprising
 - a) from 1 to 89 % by weight, preferably from 1 to 30 % by weight, of a water-soluble salen-type manganese complex,
 - b) from 10 to 95 % by weight of a dissolution restrainer,
 - c) from 0 to 20 % by weight of a further additive and
 - d) from 1 to 15 % by weight of water, based on the total weight of the granules.
2. Granules according to claim 1 that comprise as manganese complex a compound of formula



or



wherein

A is an anion;


m, n and p are each independently of the others 0, 1, 2 or 3,

R₄ is hydrogen or linear or branched C₁-C₄alkyl,


Y is a linear or branched alkylene radical of formula -[C(R₄)₂]_r-, wherein r is an integer from 1 to 8 and the R₄ radicals are each independently of the others as defined above;

-CX=CX-, wherein X is cyano, linear or branched C₁-C₈alkyl or di(linear or branched C₁-C₈alkyl)amino;

-(CH₂)_q-NR₄-(CH₂)_q-, wherein R₄ is as defined above and q is 1, 2, 3 or 4; or
a 1,2-cyclohexylene radical of formula:



or a 1,2-aryl radical of formula




wherein R₉ is hydrogen, SO₃H, CH₂OH or CH₂NH₂,

R, R₁ and R₁' are each independently of the others cyano; halogen; OR₄ or COOR₄ wherein R₄ is as defined above; nitro; linear or branched C₁-C₈alkyl; linear or branched partially fluorinated or perfluorinated C₁-C₈alkyl; or NHR₆, NR₅R₆ or N[⊖]R₅R₆R₇ wherein R₅, R₆ and R₇ are the same or different and are each hydrogen or linear or branched C₁-C₁₂alkyl or wherein R₅ and R₆ together with the nitrogen atom to which they are bonded form a 5-, 6- or 7-membered ring, which may contain further hetero atoms, or are linear or branched C₁-C₈alkyl-R₈ wherein R₈ is a radical OR₄, COOR₄ or NR₅R₆ as defined above or is NH₂ or N[⊖]R₅R₆R₇ wherein R₅, R₆ and R₇ are as defined above,

R₂ and R₃ are each independently of the other hydrogen, linear or branched C₁-C₄alkyl, unsubstituted aryl or aryl that is substituted by cyano, by halogen, by OR₄ or COOR₄ wherein

R_4 is hydrogen or linear or branched C_1 - C_4 alkyl, by nitro, by linear or branched C_1 - C_8 alkyl, by NHR_5 or NR_5R_6 , wherein R_5 and R_6 are the same or different and are each linear or branched C_1 - C_{12} alkyl or wherein R_5 and R_6 together with the nitrogen atom to which they are bonded form a 5-, 6- or 7-membered ring, which may contain further hetero atoms, by linear or branched C_1 - C_8 alkyl- R_7 wherein R_7 is an OR_4 , $COOR_4$ or NR_5R_6 radical as defined above or is NH_2 , or by $N^+(R_5R_6R_7)$ wherein R_5 , R_6 and R_7 are as defined above.

3. Granules according to claim 2 that comprise as manganese complex a compound of formula (1) or (2) wherein Y is a radical of formula $-(CH_2)_r$ wherein r is an integer from 1 to 4, especially 2, or is a radical of formula $-C(R_4)_2-(CH_2)_p-C(R_4)_2-$ wherein p is a number from 0 to 3, especially 0, and each R_4 , independently of the others, is hydrogen or C_1 - C_4 alkyl, especially hydrogen or methyl, or is a 1,2-cyclohexylene radical or a 1,2-phenylene radical of formula:



4. Granules according to either claim 2 or claim 3 that comprise as manganese complex a compound of formula (1), (2) or (3) wherein the radicals R , R_1 and R_1' are hydrogen, OR_4 , $N(R_4)_2$ or $N^+(R_4)_3$ and the R_4 groups in $N(R_4)_2$ or $N^+(R_4)_3$ may be different and are each hydrogen or C_1 - C_4 alkyl, especially methyl, ethyl or isopropyl.

5. Granules according to any one of claims 2 to 4 that comprise as manganese complex a compound of formula (1), (2) or (3) wherein the radicals R_2 and R_3 are hydrogen, methyl, ethyl or unsubstituted phenyl.

6. Granules according to any one of claims 2 to 5 that comprise as manganese complex a compound of formula (1) or (2) wherein the anion A is a halide, perchlorate, sulfate, nitrate, hydroxide, BF_4^- , PF_6^- , carboxylate, acetate, tosylate or triflate.

7. Granules according to any one of claims 2 to 6 that comprise from 1 to 30 % by weight of manganese complex of formula (1), (2) or (3), based on the total weight of the granules.

8. Granules according to any one of claims 1 to 7 that comprise as dissolution restrainer an anionic dispersing agent, a non-ionic dispersing agent or a water-soluble organic polymer.

9. Granules according to any one of claims 1 to 8 that comprise as anionic dispersing agent a condensation product of a naphthalenesulfonic acid with formaldehyde, a sodium salt of a polymerised organic sulfonic acid, a (mono-/di-)alkylnaphthalenesulfonate, a polyalkylated polynuclear arylsulfonate, a sodium salt of a polymerised alkylbenzenesulfonic acid, a lignosulfonate, an oxylignosulfonate or a condensation product of naphthalenesulfonic acid with a polychloromethyldiphenyl.

10. Granules according to any one of claims 1 to 8 that comprise as non-ionic dispersing agent a compound from the following group:

1. fatty alcohols having from 8 to 22 carbon atoms, especially cetyl alcohol,
2. addition products of preferably from 2 to 80 mol of alkylene oxide, especially ethylene oxide, in which individual ethylene oxide units may have been replaced by substituted epoxides, such as styrene oxide and/or propylene oxide, with higher unsaturated or saturated monoalcohols, fatty acids, fatty amines or fatty amides having from 8 to 22 carbon atoms, or with benzyl alcohols, phenylphenols, benzylphenols or alkylphenols in which the alkyl radicals have at least 4 carbon atoms,
3. alkylene oxide condensation products, especially propylene oxide condensation products (block polymers),
4. ethylene oxide/propylene oxide adducts with diamines, especially ethylenediamine,
5. reaction products of a fatty acid having from 8 to 22 carbon atoms with a primary or secondary amine having at least one hydroxy-lower alkyl or lower alkoxy-lower alkyl group, or alkylene oxide addition products of such hydroxyalkyl-group-containing reaction products,
6. sorbitan esters, preferably having long-chained ester groups, or ethoxylated sorbitan esters, such as, for example, polyoxyethylene-sorbitan monolaurate having from 4 to 10 ethylene oxide units or polyoxyethylene-sorbitan trioleate having from 4 to 20 ethylene oxide units,
7. addition products of propylene oxide with a tri- to hexa-hydric aliphatic alcohol having from 3 to 6 carbon atoms, for example glycerol or pentaerythritol, and

8. fatty alcohol polyglycol mixed ethers, especially addition products of from 3 to 30 mol of ethylene oxide and from 3 to 30 mol of propylene oxide with aliphatic monoalcohols having from 8 to 22 carbon atoms.

11. Granules according to claim 9 that comprise as non-ionic dispersing agent a surfactant of formula



wherein

R_{11} is C_8-C_{22} alkyl or C_8-C_{18} alkenyl;

R_{12} is hydrogen; C_1-C_4 alkyl; a cycloaliphatic radical having at least 6 carbon atoms or benzyl;

"alkylene" is an alkylene radical having from 2 to 4 carbon atoms and

n is a number from 1 to 60.

12. Granules according to any one of claims 1 to 8 that comprise as water-soluble polymer a compound from the following group:

polyethylene glycols, copolymers of ethylene oxide with propylene oxide, gelatin, polyacrylates, polymethacrylates, polyvinylpyrrolidones, vinylpyrrolidones, vinyl acetates, polyvinylimidazoles, polyvinylpyridine N-oxides, copolymers of vinylpyrrolidone with long-chained α -olefins, copolymers of vinylpyrrolidone with vinylimidazole, poly(vinylpyrrolidone/dimethylaminoethyl methacrylates), copolymers of vinylpyrrolidone/dimethylaminopropyl methacrylamides, copolymers of vinylpyrrolidone/dimethylaminopropyl acrylamides, quaternised copolymers of vinylpyrrolidones and dimethylaminoethyl methacrylates, terpolymers of vinylcaprolactam/vinylpyrrolidone/dimethylaminoethyl methacrylates, copolymers of vinylpyrrolidone and methacrylamidopropyl-trimethylammonium chloride, terpolymers of caprolactam/vinylpyrrolidone/dimethylaminoethyl methacrylates, copolymers of styrene and acrylic acid, polycarboxylic acids, polyacrylamides, carboxymethylcellulose, hydroxymethylcellulose, polyvinyl alcohols, optionally hydrolysed polyvinyl acetate, copolymers of ethyl acrylate with methacrylate and methacrylic acid, copolymers of maleic acid with unsaturated hydrocarbons and mixed polymerisation products of the said polymers.

13. Granules according to claim 12 that comprise as organic polymer carboxymethylcellulose, a polyacrylamide, a polyvinyl alcohol, a polyvinylpyrrolidone, gelatin, a hydrolysed

polyvinyl acetate, a copolymer of vinylpyrrolidone and vinyl acetate, a polyacrylate, a copolymer of ethyl acrylate with methacrylate and methacrylic acid or a polymethacrylate.

14. Granules according to claim 1 that comprise the dissolution restrainer in an amount of from 10 to 95 % by weight, preferably from 15 to 85 % by weight and especially from 25 to 75 % by weight, based on the total weight of the granules.

15. A washing agent formulation comprising

I) from 5 to 90 %, preferably from 5 to 70 %, A) of an anionic surfactant and/or B) of a non-ionic surfactant,

II) from 5 to 70 %, preferably from 5 to 50 %, especially from 5 to 40 %, C) of a builder substance,

III) from 0.1 to 30 %, preferably from 1 to 12 %, D) of a peroxide and

IV) granules according to any one of claims 1 to 14 in such an amount that the washing agent formulation comprises from 0.005 to 2 %, preferably from 0.02 to 1 %, especially from 0.1 to 0.5 %, of the pure manganese complex of formula (1), (2) or (3), the percentage figures in each case being percentages by weight based on the total weight of the washing agent.

1048045.012402

DECLARATION AND POWER OF ATTORNEY FOR U.S. PATENT APPLICATIONS

☐ Original ☐ Supplemental ☐ Substitute ☒ PCT

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if more than one name is listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Water-soluble granules of salen-type manganese complexes

which is described and claimed in:

- ☐ the attached specification.
- ☐ the specification in U.S. Application No. _____
filed _____ (day/month/year), and as amended on _____ (day/month/year) (if applicable).
- ☒ the specification in International Application No. **PCT/EP 00/06934**
filed **20 July 2000**
(day/month/year)
assigned U.S. Application No. _____ (if applicable), and as amended
☐ under PCT Article 19 on _____ (day/month/year) (if applicable)
☐ under PCT Article 34 on _____ (day/month/year) (if applicable)
☐ and further amended on _____ (day/month/year) (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information which is known by me to be material to the patentability of this application as defined in 37 C.F.R. § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119 (a)-(d) of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America relating to this subject matter having a filing date before that of the application on which priority is claimed:

COUNTRY/REGION (OR PCT)	APPLICATION No.	FILING DATE (day/month/year)	PRIORITY CLAIMED
Europe (designating DE)	99810684.3	28 July 1999	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> No

I hereby claim the benefit under 35 U.S.C. § 119 (e) of any United States provisional application(s) listed below:

APPLICATION NO.	FILING DATE (day/month/year)
-----------------	---------------------------------

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s) or PCT international application(s) designating the United States listed below and, insofar as the application discloses and claims subject matter in addition to that disclosed in the prior copending application, I acknowledge the duty to disclose all information known by me to be material to patentability as defined in 37 C.F.R. § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

U.S. APPLICATION No.	FILING DATE (day/month/year)	STATUS
<input type="checkbox"/>	Patented	<input type="checkbox"/> Pending <input type="checkbox"/> Abandoned
<input type="checkbox"/>	Patented	<input type="checkbox"/> Pending <input type="checkbox"/> Abandoned
<input type="checkbox"/>	Patented	<input type="checkbox"/> Pending <input type="checkbox"/> Abandoned
<input type="checkbox"/>	Patented	<input type="checkbox"/> Pending <input type="checkbox"/> Abandoned
<input type="checkbox"/>	Patented	<input type="checkbox"/> Pending <input type="checkbox"/> Abandoned

PCT APPLICATION No. (designating the U.S.)	INTERNATIONAL FILING DATE (day/month/year)	U.S. APPLICATION No. (if any)	STATUS
			<input type="checkbox"/> Patented <input type="checkbox"/> Pending <input type="checkbox"/> Abandoned

10043045.012402

I hereby appoint the following attorneys and agents, associated with Customer No. 000324, each of them with full power of substitution, revocation and appointment of associates, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Luther A. R. Hall (Reg. No. 27,337), JoAnn L. Villamizar (Reg. No. 30,598), Kevin T. Mansfield (Reg. No. 31,635), David R. Crichton (Reg. No. 37,300), Michele A. Kovaleski (Reg. No. 37,865) and Tyler A. Stevenson (Reg. No. 46,388).


Address all correspondence associated with Customer No. 000324 to *Ciba Specialty Chemicals Corporation, Patent Department, 540 White Plains Road, P.O. Box 2005, Tarrytown, NY 10591-9005.*

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. § 1001, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole
or first joint inventor

Menno HAZENKAMP

Inventor's signature



Date 6.12.01
(day/month/year)

Residence

Obere Weid 10
4125 Riehen
Switzerland

CHX

Citizenship

Dutch

Post Office Address

same as above

2-00
Full name of second
joint inventor, if any **Frank BACHMANN**

Inventor's signature

Frank Bachmann

Date

07.12.01
(day/month/year)

Residence

Lehener Strasse 148
79106 Freiburg
Germany *DET*

Citizenship

German

Post Office Address

same as above

3-00
Full name of third
joint inventor, if any

Cornelia MAKOWKA

Inventor's signature

Cornelia Makowka

Date

07.12.2001
(day/month/year)

Residence

Scheffelstrasse 14
79725 Laufenburg
Germany *DET*

Citizenship

German

Post Office Address

same as above

4-00
Full name of fourth
joint inventor, if any

Petr KVITA

Inventor's signature

Petr Kvita

Date

10.12.2001
(day/month/year)

Residence

Binningerstrasse 4D
4153 Reinach
Switzerland *CHX*

Citizenship

Swiss

Post Office Address

same as above

500
Full name of fifth
joint inventor, if any

Rolf KURATLI

Inventor's signature

R. Kuratli

Date

11.12.2001
(day/month/year)

Residence

Im Heimatland 8

4058 Basel

Switzerland

CHX

Citizenship

Swiss

Post Office Address

same as above

6
Full name of sixth
joint inventor, if any

Anita SCHMIDLIN

Inventor's signature

A. Schmidlin

Date

11.12.2001
(day/month/year)

Residence

Bodenackerstrasse 6C

4334 Sisseln

Switzerland

CHX

Citizenship

Swiss

Post Office Address

same as above

Full name of seventh
joint inventor, if any

Inventor's signature

Date

(day/month/year)

Residence

Citizenship

Post Office Address

same as above